

**CAUTION**

**KEEP OUT REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**SLINGSHOT™ FUNGICIDE**

**ACTIVE CONSTITUENT: 125g/L TRIADIMEFON**

GROUP	<b>C</b>	FUNGICIDE
-------	----------	-----------

For the control of fungal diseases of wheat, barley, peas, grapevines, cucurbits and sugar cane as per the Directions for Use Table.

**CONTENTS: 20L**

**SIPCAM PACIFIC AUSTRALIA PTY LIMITED**

**Suite 11, 23-31 Gheringhap Street**

**Geelong VIC 3220**

**ACN 073 176 888**

**™ Trademark of Sipcaml Pacific Australia Pty Ltd**

## DIRECTIONS FOR USE

Crop	Disease	State	Rate	WHP (days)	Critical Comments
Barley	Scald	NSW, Vic, Tas, SA, WA only	1L/ha	28	Where this disease frequently affects the flag leaf, apply at or near mid-tillering.
	Powdery mildew	All States		28	Apply at first signs of stress.
Wheat	Rust stripe (leaf and stem)	Mid and North Western Victoria	500mL or 1L/ha		For spray timings see general instructions. Normally use 500mL/ha. Only use 1L/ha where the yield is expected to be greater than 4-5 tonnes/ha or if a longer residual is required.
	Speckled leaf blotch		1L/ha		Where this disease commonly affects the Flag leaf, apply to susceptible varieties at the 5-6 tiller stage. Re-infestation can occur if moist conditions prevail.
Peas	Powdery Mildew	NSW, Vic, Tas only	500mL/ha	14	Apply as a preventative spray at flowering, at the first sign of disease, or if disease is imminent. Apply a second spray 14 days later (flat pod stage).
Grapevines	Powdery mildew	NSW, Vic, WA only	20mL/100L or 200mL/ha	14	Spray in the following method: <ol style="list-style-type: none"> <li>1. When shoots are 10cm in length.</li> <li>2. Immediately before flowering.</li> <li>3. Mid flowering.</li> <li>4. After fruit set.</li> <li>5. Prior to bunch close.</li> </ol>
Sugar Cane	Pineapple disease	Qld, NSW only	100mL in 100L	-	Fully immerse, dip or spray cut ends. Ensure thorough wetting for up 3 minutes.
Cucurbits eg melon, pumpkin, cucumber	Powdery mildew	NSW, WA only	40mL/100L or 400mL/ha	1	Apply thoroughly at the first sign of disease. Repeat at 5-10 day intervals. Use short spray intervals when disease pressure is high or plants are rapidly growing. Results are dependant on good coverage.
Azaleas	Petal blight	All States	200mL/100L	-	Apply thoroughly every 2-3 weeks from expanded bud stage until end of flowering.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

## **WITHHOLDING PERIODS:**

**Barley, Wheat:**

**DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.**

**Grapes, peas:**

**DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

**Cucurbits:**

**DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.**

**Sugar Cane, Azaleas**

**NOT REQUIRED WHEN USED AS DIRECTED**

## **GENERAL INSTRUCTIONS**

This product should be used in a full spray programme on Grapevines for prevention of Powdery Mildew infections.

This product may be used by Aircraft or Ground spraying equipment.

Spray timings for Stripe Rust control: Determine the susceptibility of the Wheat variety eg Department of Agriculture data, to determine spray requirements.

Seedling infections: When approximately 20% of leaves show signs of infection during tillering or jointing, spray within a week.

Adult infections:

***Susceptible varieties:*** Spray within one week if approximately 10% of leaves show infection. Do not delay.

***Moderately Susceptible varieties:*** Spray within 1 week if approximately 15-20% of leaves show infection. Do not delay.

***Moderately Resistant and Resistant Varieties:*** Monitor vigilantly. Spray within 1 week if rust appears and spreads.

**Mixing:** Mix the required amount of product with a small quantity of clean, filtered water to make a creamy mix. Add this blend to the required quantity of water in the tank and mix thoroughly. Agitate prior to and during spraying.

Do not mix more than can be used for the days spraying.

Do not tank mix with trace elements or other chemicals without first checking compatibility.

## **Spraying:**

**Ground Rig:** Use 50 to 100L water per hectare.

**Aerial Application** – Apply in a minimum of 10 litres of water per hectare. Apply preferably in a cross wind of greater than 5 knots. If this is not possible, increase the volume to 30L/ha and increase the droplet size. Spray in the morning and reduce the swath width. Booms with fan nozzles set forward at 45<sup>0</sup> forward to the airstream and Micronairs with blade settings of 30 – 50<sup>0</sup> are equally satisfactory. Use proper measures (eg smoke generators) for gauging wind speed, direction and variability of the wind.

## **Fungicide Resistance Warning**

GROUP	<b>C</b>	FUNGICIDE
-------	----------	-----------

Slingshot Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide.

Some naturally occurring individual fungi resistant to Slingshot Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Slingshot Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcam Pacific Australia Pty Limited accepts no liability for any losses that may result from the failure of Slingshot Fungicide to control resistant fungi.

### **Protection of Livestock**

Feed treated with this product must not be used for animal consumption, poultry feed or mixed with animal feed.

DO NOT mix leaves treated with this product with feed intended for animal consumption.

### **Protection of Wildlife, Fish, Crustaceans and Environment**

Harmful to fish. DO NOT contaminate streams, rivers or waterways with chemical or used containers.

DO NOT feed treated leaf matter to wild or domestic birds or other animals.

### **Storage and Disposal**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

### **Safety Directions**

Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear PVC or rubber apron, elbow length PVC gloves and face shield. When using prepared spray wear a face shield. If product on skin immediately wash area with soap and water. If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

### **First Aid**

If poisoning occurs contact a doctor or Poisons Information Centre(Tel: 131126)  
If swallowed, and if more than 15 minutes form a hospital, induce vomiting , preferably using  
Ipecac Syrup APF.

### **Material Safety Data Sheet**

Additional information is listed in the Material Safety Data Sheet which is available from the  
supplier.

### **NOTICE TO BUYER**

This product is not to be used for any purpose or in any way contrary to label instructions.  
The buyer by use or application of this product acknowledges so doing upon the terms and  
conditions agreed between Sipcam Pacific Pty Limited and the buyer but not limited to the  
following.

To the extent permitted by law all conditions and other warranties or other rights of action  
which buyer or any other user may have against Sipcam Pacific Pty Limited (Sipcam) or  
Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will  
not accept responsibility for any indirect or consequential loss arising from reliance on  
product information or advice provided by Sipcam or on its behalf unless it can be  
established that such information or advice was provided negligently and that the product has  
been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not  
be held liable for any loss, injury or damage indirect or consequential arising form the sale,  
supply, use or application of this product. Sipcam's liability shall in all circumstances be  
limited to replacement of the product or a refund of the purchase price paid therefor.

NRA Approval No:

Batch No:

Date of Manufacture:

THIS PRODUCT IS NOT CLASSIFIED AS A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS BY ROAD AND RAIL.
IN A TRANSPORT EMERGENCY <b>DIAL 000</b> POLICE OR FIRE BRIGADE FOR SPECIALIST ADVICE, CONTACT UNITED TRANSPORT SERVICES EMERGENCY RESPONSE COORDINATOR: <b>1800 033 111</b> (24 HOURS)